

HD CHOREA TREATMENT INFORMATION



For the treatment of **Huntington's disease (HD) chorea in adults**

Reduce HD chorea movements with once-daily AUSTEDO XR®



[#1 prescribed treatment for HD chorea*]

[*Time period 05/2022 through 04/2025.]

APPROVED USES

AUSTEDO XR® (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat:

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR® and AUSTEDO® can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Please read **Important Safety Information** on pages 8-9 and click [here](#) or visit www.AUSTEDO.com to read or print the **Medication Guide** for AUSTEDO XR.



What is Huntington's disease (HD) chorea?

SYMPTOMS AND CAUSES OF HD CHOREA

HD is a genetic brain disorder that causes an abnormal increase in the release of dopamine. Dopamine is a brain chemical responsible for movement, and increased amounts in the brain can cause chorea. Chorea is a symptom of HD that causes uncontrollable body movement.

*Chorea is one of the most visible symptoms of HD and **can be treated***

Chorea can be a burden to people with HD and those who care for them, impacting daily life and relationships. Care partners reported chorea as the most impactful symptom of HD.

What HD chorea signs look like:



Flicking the fingers



Flailing arms or legs



Shrugging shoulders



Grimacing



Twisting or jerking



Irregular eye movements or blinking

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take AUSTEDO XR or AUSTEDO if you:

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.

THE IMPACT OF HUNTINGTON'S DISEASE (HD) CHOREA

The movements of HD chorea can:



Make walking and climbing stairs difficult



Make getting dressed challenging



Affect your ability to brush teeth or bathe



Affect your ability to eat or chew



Make it difficult to speak clearly



People with chorea cannot stop these movements on their own and the movements may worsen with stress. Over time, these movements cause a loss of independence.

Learn about the **treatment benefits of AUSTEDO XR[®]**
(deutetrabenazine) extended-release tablets

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take AUSTEDO XR or AUSTEDO if you:

- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.

About treatment

BENEFITS OF AUSTEDO XR® (DEUTETRABENAZINE) EXTENDED-RELEASE TABLETS



AUSTEDO XR reduces the movements of HD chorea with **a once-daily, extended-release pill.***



People taking AUSTEDO XR reported being **more physically capable** of performing everyday tasks.



Your HD chorea treatment needs are unique to you, and AUSTEDO XR has **different dosing strengths** so your healthcare provider can find the one that works.



Scan this QR code or visit www.AUSTEDO.com to view the common side effects of AUSTEDO XR.

*Once-daily AUSTEDO XR contains the same active ingredient as twice-daily AUSTEDO® (deutetrabenazine) tablets. Data on this page is based on twice-daily dosing.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Do not take AUSTEDO XR or AUSTEDO if you:

- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

GAINING CONTROL OVER HUNTINGTON'S DISEASE (HD) CHOREA MAY BE POSSIBLE

>9^{OUT} OF 10 PEOPLE

on AUSTEDO XR® (deutetrabenazine) extended-release tablets said their movement control had improved since starting treatment†

People on AUSTEDO® saw MOVEMENT REDUCTION AS EARLY AS TWO WEEKS

ESTABLISHED SAFETY RESULTS ~3 YEARS‡



Ray, a real person living with HD chorea and Rhonda, a real HD chorea care partner.

100% of HD patients found it easy to take AUSTEDO XR†

†Results from a real-world survey including 31 patients with HD chorea taking AUSTEDO XR.

‡This was an open-label study, which is a drug trial in which all participants are aware of the drug being tested.

IMPORTANT SAFETY INFORMATION (CONTINUED)


Other possible serious side effects include:

- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.

How to get started

PERSONALIZED DOSING FOR HUNTINGTON'S DISEASE (HD) CHOREA

Your healthcare provider will start you at a low dose of AUSTEDO XR® (deutetrabenazine) extended-release tablets and gradually increase your dose over time. This approach allows your body to get used to the medication and can help with side effects as movements are reduced. This way, your healthcare provider can find a dose that works for you. This process is called titration.



Pills are for illustration only and not actual size or likeness.

Getting to your maintenance dose

Once you reach a dose that provides effective HD chorea movement reduction and is tolerated, that will become your maintenance dose. If you want to increase your dose, you and your healthcare provider can discuss more dosing options.



Because everyone is different, your maintenance dose will be the one that works specifically for you



IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.

HOW TO TAKE YOUR MEDICATION



Once daily



With water



Whole (do not
crush or break)



With or
without food

If you cannot swallow AUSTEDO XR®/AUSTEDO® (deutetrabenazine) tablets whole, tell your healthcare provider. You may need a different medication.

HOW YOU'LL BEGIN TREATMENT



Not actual image of the free Titration Kit.

Your healthcare provider may provide you a FREE Titration Kit to either:

Take home from their
office

— **OR** —

Pick up at your
pharmacy*

The Titration Kit will guide you through your first month with AUSTEDO XR.

*With free trial voucher at AUSTEDOcardform.com

97% of HD patients found it easy to include
AUSTEDO XR in their daily routine[†]

[†]Results from a real-world survey including 31 patients with HD chorea taking AUSTEDO XR.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

APPROVED USES

AUSTEDO XR® (deutetrabenazine) extended-release tablets and AUSTEDO® (deutetrabenazine) tablets are prescription medicines that are used to treat:

- the involuntary movements (chorea) of Huntington's disease. AUSTEDO XR and AUSTEDO do not cure the cause of the involuntary movements, and it does not treat other symptoms of Huntington's disease, such as problems with thinking or emotions.
- movements in the face, tongue, or other body parts that cannot be controlled (tardive dyskinesia).

It is not known if AUSTEDO XR and AUSTEDO are safe and effective in children.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR® and AUSTEDO® can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Do not take AUSTEDO XR or AUSTEDO if you:

- have Huntington's disease and are depressed or have thoughts of suicide.
- have liver problems.
- are taking reserpine. **Do not** take medicines that contain reserpine with AUSTEDO XR or AUSTEDO. If your healthcare provider plans to switch you from taking reserpine to AUSTEDO XR or AUSTEDO, you must wait at least 20 days after your last dose of reserpine before you start taking AUSTEDO XR or AUSTEDO.
- are taking a monoamine oxidase inhibitor (MAOI) medicine. **Do not** take an MAOI within 14 days after you stop taking AUSTEDO XR or AUSTEDO. **Do not** start AUSTEDO XR or AUSTEDO if you stopped taking an MAOI in the last 14 days. Ask your healthcare provider or pharmacist if you are not sure.
- are taking tetrabenazine. If your healthcare provider plans to switch you from tetrabenazine to AUSTEDO XR or AUSTEDO, take your first dose of AUSTEDO XR or AUSTEDO on the day after your last dose of tetrabenazine.
- are taking valbenazine.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Other possible serious side effects include:

- **Irregular heartbeat (QT prolongation).** AUSTEDO XR and AUSTEDO increases your chance of having certain changes in the electrical activity in your heart. These changes can lead to a dangerous abnormal heartbeat. Taking AUSTEDO XR or AUSTEDO with certain medicines may increase this chance.
- **Neuroleptic Malignant Syndrome.** Call your healthcare provider right away and go to the nearest emergency room if you develop these signs and symptoms that do not have another obvious cause: high fever, stiff muscles, problems thinking, very fast or uneven heartbeat, or increased sweating.
- **Restlessness.** You may get a condition where you feel a strong urge to move. This is called akathisia.
- **Parkinsonism.** Symptoms include: slight shaking, body stiffness, trouble moving, trouble keeping your balance, or falls.

Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO. While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

The most common side effects of AUSTEDO in people with Huntington's disease include sleepiness (sedation), diarrhea, tiredness, and dry mouth.

The most common side effects of AUSTEDO in people with tardive dyskinesia include inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).

The most common side effects of AUSTEDO XR are expected to be similar to AUSTEDO in people with Huntington's disease or tardive dyskinesia.

These are not all the possible side effects of AUSTEDO XR or AUSTEDO. Call your doctor for medical advice about side effects. You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please read the accompanying **Medication Guide** [here](#).

IT'S SIMPLE TO ENROLL IN TEVA TOTAL SUPPORT™

From benefits verification and financial assistance to nurses available to support you, we provide people with HD chorea and their care partners assistance at any point in the treatment journey.



**Starting
treatment**

**Savings and
financial support**

**Staying on
treatment**

STARTING TREATMENT

Help navigating insurance coverage

Whether you're on Medicare, privately insured, underinsured, or not insured at all, Teva **Total Support** will verify coverage and work through financial assistance options to help make treatment affordable.



**89% of people taking AUSTEDO XR®
are successfully approved for insurance coverage.**

Personalized Specialty Pharmacy support

Teva **Total Support** can help put you in touch with a pharmacy that will fulfill your prescription for AUSTEDO XR and provide ongoing assistance after you start.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Sleepiness (sedation) is a common side effect of AUSTEDO XR and AUSTEDO. While taking AUSTEDO XR or AUSTEDO, do not drive a car or operate dangerous machinery until you know how AUSTEDO XR or AUSTEDO affects you. Drinking alcohol and taking other drugs that may also cause sleepiness while you are taking AUSTEDO XR or AUSTEDO may increase any sleepiness caused by AUSTEDO XR and AUSTEDO.

Teva Total Support

SAVINGS AND FINANCIAL SUPPORT

Financial support to start and stay on treatment

New patients can start their **first 30 days free** with a **Free Trial Voucher**.*

Eligible patients may pay as little as **\$0 per month** out-of-pocket for refill prescriptions with the **AUSTEDO XR® Copay Card**.†

Medicare offers additional programs to help make treatment more affordable.

STAYING ON TREATMENT

Dedicated nurse support

Get connected with a nurse dedicated to one-on-one support throughout your treatment journey.

You can expect:

- **Outreach** calls for check-ins and goal-setting
- **Education** about your diagnosis and prescription
- **Assistance** with staying on treatment
- **Tools and resources** to help you stay on track



Additional resources are available when you enroll

Enroll in Teva Total Support

teva Total Support



1-800-887-8100

M-F, 9AM to 8PM ET

Or visit [TevaTotalSupportHD.com](https://www.TevaTotalSupportHD.com)

*The 30-day Free Trial Voucher is only for new patients. Voucher is good for up to 4 weeks of titration or 30 days of medication.

†Offer is only available to patients with commercial insurance. Offer is NOT available to patients eligible for Medicare, Medicaid, or any other government payer coverage. Out-of-pocket costs may vary. Exclusions and limitations apply. Please see complete Terms and Conditions at [AUSTEDOcardform.com](https://www.AUSTEDOcardform.com).

IMPORTANT SAFETY INFORMATION (CONTINUED)

The most common side effects of AUSTEDO in people with Huntington's disease include sleepiness (sedation), diarrhea, tiredness, and dry mouth.

The most common side effects of AUSTEDO in people with tardive dyskinesia include inflammation of the nose and throat (nasopharyngitis) and problems sleeping (insomnia).

START TREATMENT TO REDUCE HUNTINGTON'S DISEASE (HD) CHOREA

- **More than 2x movement reduction at 12 weeks vs placebo****
- **Once-daily, extended-release pill**
- **Can be taken with other medications**
- **Start at no cost**
- **Enroll in Teva Total Support for treatment benefits**



Scan this code

to access other resources. Escanee este código para acceder a otros recursos.

*Once-daily AUSTEDO XR[®] contains the same active ingredient as twice-daily AUSTEDO[®] (deutetrabenazine) tablets. Data on this page is based on twice-daily dosing.

†4.4-point improvement in people taking AUSTEDO vs 1.9 in people taking placebo based on Total Maximal Chorea (TMC) score. The TMC score is used to measure Huntington's disease (HD) chorea movements in the face, mouth, trunk, both arms, and both legs.

IMPORTANT SAFETY INFORMATION

AUSTEDO XR[®] and AUSTEDO[®] can cause serious side effects in people with Huntington's disease, including: depression, suicidal thoughts, or suicidal actions. Do not start taking AUSTEDO XR or AUSTEDO if you are depressed (have untreated depression or depression that is not well controlled by medicine) **or** have suicidal thoughts. Pay close attention to any changes, especially sudden changes, in mood, behaviors, thoughts or feelings. This is especially important when AUSTEDO XR or AUSTEDO is started and when the dose is changed. Call your healthcare provider right away if you become depressed, have unusual changes in mood or behavior, or have thoughts of suicide.

Please read **Important Safety Information** on pages 8-9 and click [here](#) or visit www.AUSTEDO.com to read or print the **Medication Guide** for AUSTEDO XR.